



COMMISSION IMPLEMENTING REGULATION (EU) 2025/839

of 5 May 2025

granting a Union authorisation for the biocidal product family ‘Oxivir Excel BPF’ in accordance with Regulation (EU) No 528/2012 of the European Parliament and of the Council

(Text with EEA relevance)

THE EUROPEAN COMMISSION,

Having regard to the Treaty on the Functioning of the European Union,

Having regard to Regulation (EU) No 528/2012 of the European Parliament and of the Council of 22 May 2012 concerning the making available on the market and use of biocidal products ⁽¹⁾, and in particular Article 44(5), first subparagraph, thereof,

Whereas:

- (1) On 16 December 2016, Diversey Europe Operations B.V. submitted an application to the European Chemicals Agency (‘the Agency’) in accordance with Article 43(1) of Regulation (EU) No 528/2012 for Union authorisation of a biocidal product family named ‘Oxivir Excel BPF’ of product-types 2 and 4, as described in Annex V to that Regulation, providing written confirmation that the competent authority of the Netherlands had agreed to evaluate the application. The application was recorded under case number BC-DL028504-45 in the Register for Biocidal Products.
- (2) ‘Oxivir Excel BPF’ contains hydrogen peroxide as the active substance, included in the Union list of approved active substances referred to in Article 9(2) of Regulation (EU) No 528/2012 for product-types 2 and 4.
- (3) On 25 September 2023, the evaluating competent authority submitted, in accordance with Article 44(1) of Regulation (EU) No 528/2012, an assessment report and the conclusions of its evaluation to the Agency.
- (4) On 20 March 2024, the Agency submitted to the Commission its opinion ⁽²⁾, the draft summary of the biocidal product characteristics (‘SPC’) of ‘Oxivir Excel BPF’ and the final assessment report on the biocidal product family, in accordance with Article 44(3) of Regulation (EU) No 528/2012.
- (5) The opinion concludes that ‘Oxivir Excel BPF’ is a biocidal product family within the meaning of Article 3(1), point (s), of Regulation (EU) No 528/2012, that it is eligible for Union authorisation in accordance with Article 42(1) of that Regulation and that, subject to compliance with the draft SPC, it meets the conditions laid down in Article 19(6) of that Regulation.
- (6) On 15 April 2024, the Agency transmitted to the Commission the draft SPC in all the official languages of the Union in accordance with Article 44(4) of Regulation (EU) No 528/2012.
- (7) The Commission concurs with the opinion of the Agency and considers it therefore appropriate to grant a Union authorisation for the biocidal product family ‘Oxivir Excel BPF’.
- (8) The measures provided for in this Regulation are in accordance with the opinion of the Standing Committee on Biocidal Products,

⁽¹⁾ OJ L 167, 27.6.2012, p. 1, ELI: <http://data.europa.eu/eli/reg/2012/528/oj>.

⁽²⁾ ECHA opinion of 29 February 2024 on the Union authorisation of ‘Oxivir Excel BPF’ (ECHA/BPC/417/2024), <https://echa.europa.eu/opinions-on-union-authorisation>.

HAS ADOPTED THIS REGULATION:

Article 1

A Union authorisation with authorisation number EU-0032483-0000 is hereby granted to Diversey Europe Operations B.V. for the making available on the market and use of the biocidal product family 'Oxivir Excel BPF' in accordance with the summary of the biocidal product characteristics set out in the Annex.

The Union authorisation is valid from 26 May 2025 until 30 April 2035.

Article 2

This Regulation shall enter into force on the twentieth day following that of its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 5 May 2025.

For the Commission
The President
Ursula VON DER LEYEN

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT FAMILY

Oxivir Excel BPF

Product type(s)

PT02: Disinfectants and algaecides not intended for direct application to humans or animals

PT04: Food and feed area

Authorisation number EU-0032483-0000**R4BP asset number** EU-0032483-0000

PART I

FIRST INFORMATION LEVEL

1. ADMINISTRATIVE INFORMATION

1.1. Family name

Name	Oxivir Excel BPF
------	------------------

1.2. Product type(s)

Product type(s)	PT02: Disinfectants and algaecides not intended for direct application to humans or animals PT04: Food and feed area
-----------------	---

1.3. Authorisation holder

Name and address of the authorisation holder	Name	Diversey Europe Operations B.V.
	Address	De Corridor 4 (Regulatory team) 3621 ZB Breukelen NL
Authorisation number	EU-0032483-0000	
R4BP asset number	EU-0032483-0000	
Date of the authorisation	26 May 2025	
Expiry date of the authorisation	30 April 2035	

1.4. Manufacturer(s) of the product

Name of manufacturer	Diversey Europe Operations B.V.
Address of manufacturer	De Corridor 4 (Regulatory team) 3621 ZB Breukelen NL
Location of manufacturing sites	

Name of manufacturer	Multifill BV
Address of manufacturer	Constructieweg 25a 3640 AJ Mijdrecht Netherlands (the)
Location of manufacturing sites	Multifill BV site 1 Constructieweg 25a 3640 AJ Mijdrecht Netherlands (the)

Name of manufacturer	Pluswipes LTD
Address of manufacturer	Pywell Rd, Willowbrook East Industrial Estate NN17 5XJ Corby United Kingdom of Great Britain and Northern Ireland (the)
Location of manufacturing sites	Pluswipes LTD site 1 Pywell Rd, Willowbrook East Industrial Estate NN17 5XJ Corby United Kingdom of Great Britain and Northern Ireland (the)

Name of manufacturer	Flexible Medical Packaging Ltd
Address of manufacturer	Unit 8, Hightown - White Cross Industrial Estate LA1 4XS Lancanster, Lancashire United Kingdom of Great Britain and Northern Ireland (the)
Location of manufacturing sites	Flexible Medical Packaging Ltd site 1 Unit 8, Hightown - White Cross Industrial Estate LA1 4XS Lancanster, Lancashire United Kingdom of Great Britain and Northern Ireland (the)

Name of manufacturer	Entegris Cleaning Process (ECP) S.A.S
Address of manufacturer	395 rue Louis Lépine 34000 Montpellier France
Location of manufacturing sites	Entegris Cleaning Process (ECP) S.A.S site 1 395 rue Louis Lépine 34000 Montpellier France

Name of manufacturer	Cheport spol s.r.o.
Address of manufacturer	Lhotsko 93 76312 Vizovice Czechia
Location of manufacturing sites	Cheport spol s.r.o. site 1 Lhotsko 93 76312 Vizovice Czechia

Name of manufacturer	Helichem
Address of manufacturer	Kozakkenberg 5 5951 DL Belfeld Netherlands (the)

Location of manufacturing sites	Helichem site 1 Kozakkenberg 5 5951 DL Belfeld Netherlands (the)
Name of manufacturer	LARA KOZMETİK HİJYENİK ÜRÜNLERİ SAN. VE DIŞ TİC. A.Ş.
Address of manufacturer	Türkgücü Organize San. Bölgesi 123.Sokak No 4/1 59580 Çorlu, Tekirdağ Türkiye
Location of manufacturing sites	LARA KOZMETİK HİJYENİK ÜRÜNLERİ SAN. VE DIŞ TİC. A.Ş. site 1 Türkgücü Organize San. Bölgesi 123.Sokak No:4/1 59580 Çorlu, Tekirdağ Türkiye
Name of manufacturer	SAPRO Temizlik Ürünleri Sanayi ve Ticaret A.Ş.
Address of manufacturer	Ortaköy Mah. İltar Bulvarı No: 27 34592 Silivri, Istanbul Türkiye
Location of manufacturing sites	SAPRO Temizlik Ürünleri Sanayi ve Ticaret A.Ş. site 1 Ortaköy Mah. İltar Bulvarı No: 27 34592 Silivri, Istanbul Türkiye
Name of manufacturer	ITW Contamination Control BV
Address of manufacturer	Saffierlaan 5 2132 VZ Hoofddorp Netherlands (the)
Location of manufacturing sites	ITW Contamination Control BV site 1 Saffierlaan 5 2132 VZ Hoofddorp Netherlands (the)
Name of manufacturer	Arkema
Address of manufacturer	Tankhoofd 10 Haven 3255 3196 KE Rotterdam Netherlands (the)
Location of manufacturing sites	Arkema site 1 Morschheimer Strasse 19 D-67292 Kirchheimbolanden Germany

1.5. **Manufacturer(s) of the active substance(s)**

Active substance	Hydrogen peroxide
Name of manufacturer	Evonik Operations GmbH
Address of manufacturer	Evonik Resource Efficiency GmbH, Rellinghauser Strasse 1-11 D-45128 Essen Germany
Location of manufacturing sites	Evonik Resource Efficiency GmbH site 1 Untere Kanalstr. 3 79618 Rheinfelden Germany

Active substance	Hydrogen peroxide
Name of manufacturer	Solvay Chemicals International SA
Address of manufacturer	Solvay Chemicals International SA, Rue Ransbeek 310 1120 Brussels Belgium
Location of manufacturing sites	Solvay Chemicals International site 1 Via Piave 6 57013 Rosignano Solvay Italy Solvay Chemicals International site 2 Köthensche Strasse 1-3 06406 Bernburg Germany Solvay Chemicals International site 3 Baronet Road WA4 6HB Warrington United Kingdom of Great Britain and Northern Ireland (the) Solvay Chemicals International site 4 Yrjonojantie 2 45911 Voikkaa Finland Solvay Chemicals International site 5 Rua Eng. Clement Dumoulin, 2625-106 Povia de Santa Iria Portugal

Active substance	Hydrogen peroxide
Name of manufacturer	Kemira Oyj
Address of manufacturer	Kemira Oyj, Porkkalankatu 3, PL 330 00101 Helsinki Finland
Location of manufacturing sites	Kemira Oyj site 1 Moezelweg 151 3198 LS Rotterdam Netherlands (the) Kemira Oyj site 2 Typpitie PL 171 90101 Oulu Finland Kemira Oyj site 3 Industrigatan 83 25109 Helsingborg Sweden

Active substance	Hydrogen peroxide
Name of manufacturer	Arkema France
Address of manufacturer	420, rue d'Étienne d'Orves F-92705 Colombes Cedex France
Location of manufacturing sites	Arkema b.v. site 1 Route National 85 38560 Jarrie France Arkema b.v. site 2 Am Haupttor, Bau 2410 06237 Leuna Germany

Active substance	Hydrogen peroxide
Name of manufacturer	Arkema GmbH
Address of manufacturer	Tersteegenstr. 28 D-40474 Düsseldorf Germany

Location of manufacturing sites	Arkema b.v site 1 Route National 85 3856 Jarrie France Arkema b.v. site 2 Am Haupttor, Bau 2410 0623 Leuna Germany
---------------------------------	---

Active substance	Hydrogen peroxide
Name of manufacturer	Evonik Antwerpen NV
Address of manufacturer	Tijsmanstunnel West 2040 Antwerpen Belgium
Location of manufacturing sites	Evonik Antwerpen site 1 Tijsmanstunnel West 2040 Antwerpen Belgium

Active substance	Hydrogen peroxide
Name of manufacturer	Evonik Peroxid GmbH
Address of manufacturer	Industriestraße 1 9721 Weißenstein Austria
Location of manufacturing sites	Evonik Peroxid GmbH site 1 Industriestrasse 1 9721 Weissenstein Austria

Active substance	Hydrogen peroxide
Name of manufacturer	Evonik Peroxide Netherlands BV
Address of manufacturer	Hettenheувelweg 37 /39 1101 BM Amsterdam Netherlands (the)
Location of manufacturing sites	Evonik Peroxide Netherlands BV site 1 Oosterhorn 14 9936 HD Farmsum Netherlands (the)

Active substance	Hydrogen peroxide
Name of manufacturer	Evonik Peroxide Spain, S.L.U.
Address of manufacturer	C/ Afueras s/h. 50784 La Zaida Spain
Location of manufacturing sites	Evonik Peroxide Spain, S.L.U. site 1 Calle Afueras s/n 50784 La Zaida Spain

2. PRODUCT FAMILY COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the family

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Hydrogen peroxide		active substance	7722-84-1	231-765-0	0,36 - 7,2 % (w/w)

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
(C9-C11) Alkyl alcohol ethoxylate	(C9-C11) Alkyl alcohol ethoxylate	Non-Active substance	68439-46-3		0,09 - 1,8 % (w/w)
alkyl benzene sulphonic acid	2-Dodecylbenzenesulfonic acid	Non-Active substance	85536-14-7	287-494-3	0,97 - 19,4 % (w/w)
methane sulphonic acid		Non-Active substance	75-75-2	200-898-6	0,12 - 2,4 % (w/w)

2.2. Type(s) of formulation

Formulation type(s)	AL Any other liquid SL Soluble concentrate AL (Any other liquid) - RTU wipe
---------------------	---

PART II

SECOND INFORMATION LEVEL - META SPC(S)

1. META SPC 1 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 1 identifier

Identifier	Meta SPC: Meta-SPC 1: Concentrate
------------	-----------------------------------

1.2. Suffix to the authorisation number

Number	1-1
--------	-----

1.3. Product type(s)

Product type(s)	PT02: Disinfectants and algacides not intended for direct application to humans or animals PT04: Food and feed area
-----------------	--

2. META SPC 1 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 1

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Hydrogen peroxide		active substance	7722-84-1	231-765-0	7,2 - 7,2 % (w/w)

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
alkyl benzene sulphonic acid	2-Dodecylbenzenesulfonic acid	Non-Active substance	85536-14-7	287-494-3	19,4 - 19,4 % (w/w)
(C9-C11) Alkyl alcohol ethoxylate	(C9-C11) Alkyl alcohol ethoxylate	Non-Active substance	68439-46-3		1,8 - 1,8 % (w/w)
methane sulphonic acid		Non-Active substance	75-75-2	200-898-6	2,4 - 2,4 % (w/w)

2.2. Type(s) of formulation of the meta SPC 1

Formulation type(s)	SL Soluble concentrate
---------------------	------------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 1

Hazard statements	H314: Causes severe skin burns and eye damage. H412: Harmful to aquatic life with long lasting effects. H290: May be corrosive to metals.
Precautionary statements	P234: Keep only in original packaging. P273: Avoid release to the environment. P280: Wear protective gloves, protective clothing and eye or face protection. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower]. P310: Immediately call a POISON CENTRE, doctor or physician. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P310: Immediately call a POISON CENTRE, doctor or physician. P301+P330+P331: IF SWALLOWED: rinse mouth. Do NOT induce vomiting. P310: Immediately call a POISON CENTRE, doctor or physician. P363: Wash contaminated clothing before reuse. P390: Absorb spillage to prevent material damage. P501: Dispose of contents to hazardous waste disposal service in accordance with the statutory regulations. P501: Dispose of container to hazardous waste disposal service in accordance with the statutory regulations.

4. AUTHORISED USE(S) OF THE META SPC

4.1. Use description

Table 1.

Use # 1.1 – Cleaning and disinfection by mopping for healthcare areas.

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for hard surfaces for healthcare areas.</p>
Application method(s)	<p>Method: Mopping</p> <p>Detailed description: Mopping with a soaked mop.</p>
Application rate(s) and frequency	<p>Application rate: 11 to 40 ml/m² Dilute the product with water to reach 4,5 % product</p> <p>Number and timing of application: Frequency of use: 7 times a day</p> <p>Use at room temperature</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres</p> <p>HDPE bottle (J-Flex): 1,5 litres</p> <p>HDPE bottle (Smart Dose): 1,4 litres – 1,5 litres</p> <p>HDPE intermediate bulk container (IBC): 100-1 000 litres</p>

4.1.1. *Use-specific instructions*

Product dilution:

bactericidal, yeasticidal and virucidal 4,5% product.

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of the concentrated product to reach the dilution of 4,5%. Carefully fill up with the rest of the water.

Add the concentrated product to the water to avoid splashes of concentrated product.

Dip the mop into the container. Squeeze the surplus solution. Clean and disinfect the floor with the wet mop.

Make sure to wet the surface completely.

Allow to act for 5 minutes. Let it air dry. No rinsing needed after disinfection.

For packs with (semi-)automatic dilution system with 2 settings:

Select the right setting on the dosing equipment. Fill a container with the diluted product (4,5%). Dip the mop into the container. Squeeze the surplus solution. Clean and disinfect the floor with the wet mop. Make sure to wet the surface completely. Allow to act for 5 minutes.

Let it air dry. No rinsing needed after disinfection.

4.1.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 1

4.1.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 1

4.1.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 1

4.1.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

4.2. Use description

Table 2.

Use # 1.2– Cleaning and disinfection with a cloth or sponge for healthcare areas.

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for hard surfaces for healthcare areas</p>
Application method(s)	<p>Method: spreading</p> <p>Detailed description: Spreading with a soaked cloth or sponge.</p>
Application rate(s) and frequency	<p>Application rate: 11 to 40 ml/m² Dilute the product with water to reach 4,5% product</p> <p>Number and timing of application: Frequency of use: 7 times a day</p> <p>Use at room temperature</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2.5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres</p> <p>HDPE bottle (J-Flex): 1,5 litres</p> <p>HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres</p> <p>HDPE IBC: 100-1 000 litres</p>

4.2.1. Use-specific instructions

Product dilutions:

Bactericidal, yeasticidal & virucidal 4,5%

For packs with manual dilution system

To prevent foaming, fill a container three quarters full of water. Add the correct amount of product to reach the dilution of 4,5%. Carefully fill up with the rest of the water.

Soak the cloth or sponge in the solution. Distribute the liquid on the surface. Make sure to wet the surface completely.

Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

For packs with (semi-)automatic dilution system

Select the right setting on the dosing equipment. Fill a container with the diluted product (4,5%). Soak the cloth or sponge in the solution. Distribute the liquid on the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

4.2.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.2.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.2.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.2.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 1

4.3. Use description

Table 3.

Use # 1.3 – Cleaning and disinfection of small objects by soaking or dipping for healthcare areas.

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for small objects (e.g. scissors and brushes) for healthcare areas. The biocidal use excludes disinfection of medical devices.
Application method(s)	Method: Soaking or dipping Detailed description: Soaking or dipping in the solution.

Application rate(s) and frequency	Application rate: up to 3 litres Dilute the product with water to reach 4,5% product Number and timing of application: Frequency of use: 1 time a day Use at room temperature
Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres

4.3.1. Use-specific instructions

Product dilutions:

bactericidal, yeasticidal & virucidal 4,5%

For packs with manual dilution system:

To prevent foaming, fill a container three quarters with water. Add the correct amount of product to reach the dilution of 4,5%. Carefully fill up with the rest of the water.

Add the product to the water to avoid splashes of concentrated product.

Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

For packs with (semi-)automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with the diluted product (4,5%).

Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

The solutions for soaking or dipping can be used only once.

4.3.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.3.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.3.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.3.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 1

4.4. Use description

Table 4.

Use # 1.4 – Cleaning and disinfection by foaming for healthcare areas.

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for hard surfaces for healthcare areas.</p>
Application method(s)	<p>Method: spraying</p> <p>Detailed description: Spraying using a trigger foam. Optionally wipe to spread the solution.</p>
Application rate(s) and frequency	<p>Application rate: 11 ml/m² Dilute the product with water to reach 4,5% product</p> <p>Number and timing of application:</p> <p>Frequency of use: For use in toilets and bathrooms: 4 times a day For use in hospitals and medical practices: 8 times a day For use in laboratories: 10 times a day For use in cleanrooms: 10 times a day</p> <p>Use at room temperature</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres</p> <p>HDPE bottle (J-Flex): 1,5 litres</p> <p>HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres</p> <p>HDPE IBC: 100-1 000 litres</p>

4.4.1. Use-specific instructions

Product dilutions:

bactericidal, yeasticidal & virucidal 4,5%

For packs with manual dilution system:

To prevent foaming, fill a foaming spray bottle three quarters full of water. Add the correct amount of concentrated product. to reach the dilution of 4,5%. Carefully fill up with the rest of the water.

Apply the diluted product. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surface disinfection.

For packs with (semi-)automatic dilution system:

Select the right setting on the dosing equipment. Fill a foaming spray bottle with the diluted product. Apply the diluted product. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surface disinfection.

4.4.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 1

4.4.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 1

4.4.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 1

4.4.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

4.5. **Use description**

Table 5.

Use # 1.5 –Cleaning and disinfection using moist wipes (incl. healthcare areas)

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for hard surfaces.

Application method(s)	Method: Wiping Detailed description: Wiping with prepared moist wipes.
Application rate(s) and frequency	Application rate: 1,5 m ² per wipe Dilute the product with water to reach 5% product. Number and timing of application: Frequency of use: 8 times a day Use at room temperature
Category(ies) of users	professional
Pack sizes and packaging material	Liquid: High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres Dry Wipes are packed in a HDPE bucket with 125 wipes (polypropylene (PP) sold separately).

4.5.1. Use-specific instructions

Prepare 1,5 litres dilution by adding 75 ml Oxivir Excel to 1,425 litres water.

Apply 1,5 litres diluted product to the Dry Wipes bucket. Distribute the liquid evenly. Close the bucket.

Open the lid. Take a single wipe. Close the lid after taking a wipe. Only use wet wipes. Wipe the surface and make sure to wet the surface completely.

Allow to act for 5 minutes. Let it air dry. No rinsing needed after disinfection.

4.5.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.5.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.5.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.5.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 1

4.6. Use description

Table 6.

Use # 1.6 – Cleaning and disinfection of food contact surfaces using a mop

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for hard surfaces that get into contact with food</p>
Application method(s)	<p>Method: Mopping</p> <p>Detailed description: Mopping with a soaked mop.</p>
Application rate(s) and frequency	<p>Application rate: 11 to 40 ml/m² Dilute the product with water to reach 3,75% product</p> <p>Number and timing of application: Frequency of use: 7 times a day Use at room temperature</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres</p> <p>HDPE bottle (J-Flex): 1,5 litres</p> <p>HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres</p> <p>HDPE IBC: 100-1 000 litres</p>

4.6.1. Use-specific instructions

Product dilution: bactericidal, yeasticidal & virucidal 3,75%

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of the concentrated product to reach the dilution of 3,75%. Carefully fill up with the rest of the water.

Add the concentrated product to the water to avoid splashes of the concentrated product.

Dip the mop into the container. Squeeze the surplus solution. Clean and disinfect the floor with the wet mop.

Make sure to wet the surface completely. Allow to act for 5 minutes. Let it air dry. No rinsing needed after disinfection.

For packs with (semi-) automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with the diluted product (3,75%). Soak the mop in the solution. Distribute the liquid.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

4.6.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 1

4.6.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 1

4.6.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 1

4.6.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

4.7. **Use description**

Table 7.

Use # 1.7 – Cleaning and disinfection of food contact surfaces using a cloth or sponge.

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for hard surfaces that get into contact with food.</p>
Application method(s)	<p>Method: spreading</p> <p>Detailed description: Spreading with a soaked cloth or sponge.</p>
Application rate(s) and frequency	<p>Application rate: 11 to 40 ml/m² Dilute the product with water to reach 3,75% product</p> <p>Number and timing of application: Frequency of use: 7 times a day Use at room temperature</p>

Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres

4.7.1. Use-specific instructions

Product dilution: bactericidal, yeasticidal & virucidal 3,75%

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of the concentrated product to reach the dilution of 3,75%. Carefully fill up with the rest of the water.

Add the concentrated product to the water to avoid splashes of concentrated product.

Soak the cloth or sponge in the solution. Distribute the liquid.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

For packs with (semi-) automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with the diluted product (3,75%). Soak the cloth or sponge in the solution. Distribute the liquid.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

4.7.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.7.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.7.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.7.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 1

4.8. Use description

Table 8.

Use #1.8 – Cleaning and disinfection of food contact small objects by soaking or dipping.

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for small objects (e.g. kitchen utensils) that get into contact with food.</p>
Application method(s)	<p>Method: Soaking or dipping</p> <p>Detailed description: Soaking or dipping in the solution.</p>
Application rate(s) and frequency	<p>Application rate: up to 3 litres Dilute the product with water to reach 3,75% product</p> <p>Number and timing of application: Use at room temperature Frequency of use: 1 time a day</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres</p> <p>HDPE bottle (J-Flex): 1,5 litres</p> <p>HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres</p> <p>HDPE IBC: 100-1 000 litres</p>

4.8.1. Use-specific instructions

Product dilution: bactericidal, yeasticidal & virucidal 3,75%

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of the concentrated product. to reach the dilution of 3,75%. Carefully fill up with the rest of the water.

Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

For packs with(semi-) automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with diluted product (3,75%). Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

The solutions for soaking or dipping can be used only once.

4.8.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 1

4.8.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 1

4.8.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 1

4.8.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

4.9. **Use description**

Table 9.

Use # 1.9 – Cleaning and disinfection of food contact surfaces by foaming.

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for hard surfaces that get into contact with food.
Application method(s)	Method: spraying Detailed description: Spraying using a trigger foam. Optionally wipe to spread the solution.

Application rate(s) and frequency	Application rate: 11 ml/m ² Dilute the product with water to reach 3,75% product Number and timing of application: Use at room temperature Frequency of use: For use in kitchens and canteen: 4 times a day For use in food processing industries: 4 times a day For use on a counter in a shop/supermarket: 8 times a day
Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres

4.9.1. Use-specific instructions

Product dilution: 3,75%

For packs with manual dilution system:

To prevent foaming, fill a foaming spray bottle three quarters with full of water. Add the correct amount of the concentrated product. to reach a dilution of 3,75%. Carefully fill up with the rest of the water.

Apply the diluted product. Optionally wipe to spread the solution. Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surface disinfection.

For packs with full (semi-) automatic dilution system:

Select the right setting on the dosing equipment. Fill a foaming spray bottle with the diluted product (3,75%). Apply the diluted product. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surface disinfection.

4.9.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.9.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.9.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.9.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 1

4.10. Use description

Table 10.

Use # 1.10 –Cleaning and disinfection of food contact surfaces with moist wipes

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for hard surfaces that get in contact with food.</p>
Application method(s)	<p>Method: Wiping</p> <p>Detailed description: Wiping with prepared moist wipes.</p>
Application rate(s) and frequency	<p>Application rate: 1,5 m² per wipe Dilute the product with water to reach 5% product.</p> <p>Number and timing of application: Frequency of use: 8 times a day Use at room temperature</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>Oxivir Excel: High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres Dry Wipes are packed in a HDPE bucket with 125 wipes (PP). (sold separately)</p>

4.10.1. Use-specific instructions

Product dilution 5%

Prepare 1,5 litres dilution by adding 75 ml of the concentrated product to 1,425 litres of water. To prevent foaming, fill a container three quarters full of water.

Apply 1,5 litres of the diluted product to the Dry Wipes bucket. Distribute the liquid evenly. Close the bucket.

Open the lid, take a single wipe, close the lid after taking a wipe. Only use wet wipes. Wipe the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Let it air dry. No rinsing needed after disinfection.

4.10.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.10.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.10.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.10.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 1

4.11. Use description

Table 11.

Use # 1.11 – Cleaning and disinfection of hard surfaces by mopping in non-health care areas.

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for hard surfaces.
Application method(s)	Method: Mopping Detailed description: Mopping with a soaked mop.
Application rate(s) and frequency	Application rate: 11 to 40 ml/m ² Dilute the product with water to reach the following concentration of the product Bactericidal – 1,5% product Yeasticidal – 3,75% product Virucidal – 2% product

	Number and timing of application: Frequency of use: 7 times a day Use at room temperature
Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres

4.11.1. Use-specific instructions

Product dilutions:

Bactericidal – 1,5% product

Yeasticidal – 3,75% product

Virucidal – 2% product

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of the concentrated product to reach the appropriate dilution of the product. Carefully fill up with the rest of the water.

Add the concentrated product to the water to avoid splashes of the concentrated product. Dip the mop into the container. Squeeze the surplus solution. Clean and disinfect the floor with the wet mop. Make sure to wet the surface completely. Allow to act for 5 minutes.

Let it air dry. No rinsing needed after disinfection.

For packs with (semi-) automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with the diluted product. Dip the mop into the container. Squeeze the surplus solution. Clean and disinfect the floor with the wet mop. Make sure to wet the surface completely. Allow to act for 5 minutes.

Let it air dry. No rinsing needed after disinfection.

4.11.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.11.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.11.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.11.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 1

4.12. Use description

Table 12.

Use # 1.12 – Cleaning and disinfection of hard surfaces with a cloth or sponge in non-health care areas.

Product type	PT02: Disinfectants and algacides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for hard surfaces
Application method(s)	Method: spreading Detailed description: Spreading with a soaked cloth or sponge.
Application rate(s) and frequency	Application rate: 11 to 40 ml/m ² Dilute the product with water to reach the following concentration of the product bactericidal – 1,5% product yeasticidal – 3,75% product virucidal – 2% product Number and timing of application: Use at room temperature Frequency of use: 7 times a day
Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres

4.12.1. Use-specific instructions

Product dilutions:

bactericidal 1,5%; yeasticidal 3,75%; virucidal 2%

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of the concentrated product to reach the appropriate dilution of the product. Carefully fill up with the rest of the water.

Soak the cloth or sponge in the solution. Distribute the liquid. Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

For packs with(semi-) automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with the diluted product. Soak the cloth or sponge in the solution. Distribute the liquid.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

4.12.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 1

4.12.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 1

4.12.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 1

4.12.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

4.13. **Use description**

Table 13.

Use # 1.13 – Cleaning and disinfection of small objects by soaking or dipping in non-health care areas

Product type	PT02: Disinfectants and algacides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for small objects (e.g. scissors and brushes).
Application method(s)	Method: Soaking or dipping Detailed description: Soaking or dipping in the solution.

Application rate(s) and frequency	Application rate: up to 3 litres Dilute the product with water to reach the following concentration of the product Bactericidal – 1,5% product. Yeasticidal – 3,75% product. Virucidal – 2% product Number and timing of application: Frequency of use: 1 time a day Use at room temperature
Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres

4.13.1. Use-specific instructions

Product dilution:

Bactericidal – 1,5% product.

Yeasticidal – 3,75% product.

Virucidal – 2% product

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of the concentrated product to reach the appropriate dilution of the product. Carefully fill up with the rest of the water. Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

For packs with (semi-) automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with the diluted product.

Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

The solutions for soaking or dipping can be used only once.

4.13.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 1

4.13.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 1

4.13.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 1

4.13.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

4.14. **Use description**

Table 14.

Use # 1.14 – Cleaning and disinfection of surfaces by foaming in non-health care areas.

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Cleaning and disinfection product for hard surfaces.</p>
Application method(s)	<p>Method: spraying</p> <p>Detailed description: Spraying using a trigger foam. Optionally wipe to spread out the solution.</p>
Application rate(s) and frequency	<p>Application rate: 11 ml/m² Dilute the product with water to reach the following concentration of the product Bactericidal - 1,5% product, Yeasticidal - 3,75% product, Virucidal - 2% product</p> <p>Number and timing of application:</p> <p>Use at room temperature</p> <p>Frequency of use: If product is intended for use in toilets and bathrooms: 4 times a day If product is intended for use in cosmetic/pharmaceutical manufacturing facilities: 8 times a day If product is intended for use in laboratories: 10 times a day If product is intended for use in cleanrooms: 10 times a day</p>

Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres HDPE IBC: 100-1 000 litres

4.14.1. *Use-specific instructions*

Product dilutions:

bactericidal 1.5%; yeasticidal 3.75%; virucidal 2%

For packs with manual dilution system:

To prevent foaming, fill a foaming spray bottle three quarters full of water. Add the correct amount of the concentrated product to reach the appropriate dilution of the product. Carefully fill up with the rest of the water.

Apply the diluted product. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surfaces disinfection.

For packs with (semi) automatic dilution system:

Select the right setting on the dosing equipment. Fill a foaming spray bottle with the diluted product. Apply the diluted product. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surfaces disinfection.

4.14.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 1

4.14.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 1

4.14.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 1

4.14.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

4.15. Use description

Table 15.

Use # 1.15 – Disinfection of pre-cleaned hard surfaces.

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Disinfection product for pre-cleaned hard surfaces (incl. healthcare areas, pharmaceutical industry, cosmetic industry, laboratories and cleanrooms)</p>
Application method(s)	<p>Method: Spreading, dipping and foam spraying</p> <p>Detailed description: Spreading using a soaked mop. Spreading using a soaked cloth or sponge. Dipping in a solution. Foam spraying, optionally wipe to spread the solution</p>
Application rate(s) and frequency	<p>Application rate: Application using a soaked mop: 11 to 40 ml/m², application using a soaked cloth or sponge: 11 to 40 ml/m², dipping in a solution: 3 litres, foam spraying: 11 ml/m². Dilute the product with water to reach 2,5%</p> <p>Number and timing of application: Frequency of use: Application using a soaked mop: 7 times a day Application using a soaked cloth/sponge: 7 times a day Dipping: 1 time a day Foam spraying: bathrooms & toilets 4 times a day healthcare 8 times a day laboratories & cleanrooms 10 times a day Use at room temperature</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres – 2 litres – 2,5 litres – 3 litres – 5 litres – 10 litres – 20 litres – 50 litres – 100 litres – 200 litres</p> <p>HDPE bottle (J-Flex): 1,5 litres</p> <p>HDPE Bottle (Smart Dose): 1,4 litres – 1,5 litres</p> <p>HDPE IBC: 100-1 000 litres</p>

4.15.1. *Use-specific instructions*

Product dilutions:

Bactericidal, yeasticidal & virucidal 2,5%

For packs with manual dilution system:

To prevent foaming, fill a container three quarters full of water. Add the correct amount of concentrated product to reach the dilution of 2,5%. Carefully fill up with the rest of the water.

For packs with (semi-)automatic dilution system:

Select the right setting on the dosing equipment. Fill a container with the diluted product (2,5%).

Application using a soaked mop: Dip the mop into the container. Squeeze the surplus solution. Apply onto floor.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Application using a soaked cloth/sponge: Soak the cloth or sponge in the solution. Distribute the liquid on the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Dipping: Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes. The solutions for soaking or dipping can be used only once.

Foam spraying: Fill a spray bottle with foam head. Apply the diluted product. Optionally wipe to spread the solution. Make sure to wet the surface completely. Allow to act for 5 minutes. Only for small surfaces disinfection.

Wipe dry or let air dry. No rinsing needed after disinfection

4.15.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 1

4.15.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 1

4.15.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 1

4.15.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 1

5. GENERAL DIRECTIONS FOR USE OF THE META SPC 1

5.1. Instructions for use

For professional use:

Keep away from food, drink and animal feeding stuffs.

Wash hands after handling.

Keep only in original container.

Do not mix with other products.

5.2. Risk mitigation measures

The following personal protective equipment (PPE) is stipulated without prejudice to the application by employers of Council Directive 98/24/EC and other Union legislation in the area of health and safety at work.

The use of eye protection (consistent with the European Standard EN 166)16 321 or equivalent during handling of the concentrated product is mandatory.

Wear protective chemical resistant gloves (classified under the European Standard EN ISO 374) or equivalent during product handling phase (glove material to be specified by the authorisation holder within the product information).

Wear chemical-resistant clothing (consistent with the European Standard EN 14605) or equivalent and wear suitable protective footwear (consistent with the European Standards EN 13832) or equivalent when handling concentrated products.

See section 6 for the full titles of the EN standards.

Keep pets away from treated surfaces until dried.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid:

IF INHALED: Move to fresh air and keep at rest in a position comfortable for breathing. If symptoms: Call 112/ambulance for medical assistance. If no symptoms: Call a POISON CENTRE or a doctor.

IF SWALLOWED: Immediately rinse mouth. Give something to drink, if exposed person is able to swallow. Do NOT induce vomiting. Call 112/ambulance for medical assistance.

IF ON SKIN: Immediately wash skin with plenty of water. Thereafter take off all contaminated clothing and wash it before reuse. Continue to wash the skin with water for 15 minutes. Call a POISON CENTRE or a doctor.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing for at least 15 minutes. Call 112/ambulance for medical assistance.

Information to Healthcare personnel/doctor:

The eyes should also be rinsed repeatedly on the way to the doctor if eye exposure to alkaline chemicals (pH > 11), amines and acids like acetic acid, formic acid or propionic acid

Environmental precautions:

Do not allow to enter drainage system, surface or groundwater. Do not allow to enter the ground/soil. Dilute with plenty of water. Inform responsible authorities in case undiluted product reaches drainage system, surface or ground water or the ground/soil.

5.4. Instructions for safe disposal of the product and its packaging

Dispose of contents/container to a hazardous waste disposal service in accordance with the statutory regulations.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

2 year shelf life.

Store in the dark. Do not store at temperatures above 30°C.

6. OTHER INFORMATION

The products of this meta-SPC may also be intended for the disinfecting of Medical Device surfaces in medical and clinical areas. In such a case, their placing on the market or putting into service must comply with Regulation (EU) 2017/745 of the European Parliament and of the Council of 5 April 2017 on medical devices, amending Directive 2001/83/EC, Regulation (EC) No 178/2002 and Regulation (EC) No 1223/2009 and repealing Council Directives 90/385/EEC and 93/42/EEC (OJ L 117, 05/05/2017, p. 1) and they are medical devices classified as class IIa. Those uses are out of the scope of this biocidal registration.product family authorisation and are not covered by it.

With respect to the “Category(ies) of users” note: “Professionals (including industrial users) means trained professionals if this is required by national legislation.

The full titles of the EN standards referenced in section 5.2 “Risk mitigation measures” are:

EN 13832 - Footwear protecting against chemicals

EN ISO 374 - Protective gloves against dangerous chemicals and micro-organisms

EN ISO 16321 - Eye and face protection for occupational use

EN 14605 - Protective clothing against liquid chemicals - Performance requirements for clothing with liquid-tight (Type 3) or spray-tight (Type 4) connections, including items providing protection to parts of the body only (Types PB [3] and PB [4]).

Council Directive 98/24/EC of 7 April 1998 on the protection of the health and safety of workers from the risks related to chemical agents at work (fourteenth individual Directive within the meaning of Article 16(1) of Directive 89/391/EEC) (OJ L 131, 5.5.1998, p. 11).

7. THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 1

7.1. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)	Oxivir Excel	Market area: EU
	Oxivir Excel H+	Market area: EU
	Oxivir Excel J-flex	Market area: EU
	Oxivir Excel Smart- Dose	Market area: EU

	Oxivir Excel SD	Market area: EU
	Oxivir Excel Conc	Market area: EU
	Oxivir Excel Concen- trate	Market area: EU
	Oxivir Excel IBC	Market area: EU
	Oxivir Excel Acti	Market area: EU
	Oxivir Excel Des	Market area: EU
	Oxivir Excel Duo	Market area: EU
	Clinell H2O2 concen- trate	Market area: EU
	Antibac Oxivir concen- trate	Market area: EU
	Divosan AHP VS70	Market area: EU
	TASKI Sani AHP- Des	Market area: EU
	TASKI Sprint AHP-Des	Market area: EU
	Suma AHP D10.7	Market area: EU
	Suma AHP	Market area: EU
	Suma San AHP D10.7	Market area: EU
	Suma San AHP	Market area: EU

	ClearKlens AHP Conc	Market area: EU
	ClearKlens Excel Conc	Market area: EU
	DOMES- TOS Expert Conc	Market area: EU
	GLORIX Expert Conc	Market area: EU
	KLINEX Expert Conc	Market area: EU
	LYSO- FORM Expert Conc	Market area: EU
	SAVO Expert Conc	Market area: EU
	CIF Expert Conc	Market area: EU
	VISS Expert Conc	Market area: EU
	JIF Expert Conc	Market area: EU
	FLORAS- ZEPT Expert Conc	Market area: EU
	ANDY Expert Conc	Market area: EU
	SUPER- ANDY Expert Conc	Market area: EU
	LIFEBUOY Expert Conc	Market area: EU

			ANDY-SAN Expert Conc	Market area: EU		
Authorisation number			EU-0032483-0001 1-1			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
Hydrogen peroxide		active substance	7722-84-1	231-765-0	7,2 % (w/w)	
alkyl benzene sulphonic acid	2-Dodecylbenzenesulfonic acid	Non-Active substance	85536-14-7	287-494-3	19,4 % (w/w)	
(C9-C11) Alkyl alcohol ethoxylate	(C9-C11) Alkyl alcohol ethoxylate	Non-Active substance	68439-46-3		1,8 % (w/w)	
methane sulphonic acid		Non-Active substance	75-75-2	200-898-6	2,4 % (w/w)	

1. META SPC 2 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 2 identifier

Identifier	Meta SPC: Meta-SPC 2: RTU
------------	---------------------------

1.2. Suffix to the authorisation number

Number	1-2
--------	-----

1.3. Product type(s)

Product type(s)	PT02: Disinfectants and algacides not intended for direct application to humans or animals PT04: Food and feed area
-----------------	--

2. META SPC 2 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 2

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Hydrogen peroxide		active substance	7722-84-1	231-765-0	0,36 - 0,36 % (w/w)

2.2. **Type(s) of formulation of the meta SPC 2**

Formulation type(s)	AL Any other liquid
---------------------	---------------------

3. **HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 2**

Hazard statements	H290: May be corrosive to metals.
Precautionary statements	P102: Keep out of reach of children. P234: Keep only in original packaging. P390: Absorb spillage to prevent material damage.

4. **AUTHORISED USE(S) OF THE META SPC**4.1. **Use description**

Table 1.

Use # 2.1 – Cleaning and disinfection by mopping (incl. healthcare areas).

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for hard surfaces (incl. healthcare areas)
Application method(s)	Method: Mopping Detailed description: Mopping with a soaked mop.
Application rate(s) and frequency	Application rate: 11 to 40 ml/m ² Ready to use product Number and timing of application: Use at room temperature Frequency of use: 7 times a day

Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	Professional and non-professional users: High Density Polyethene (HDPE)Canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres

4.1.1. Use-specific instructions

Fill a container with the product. Dip the mop into the container. Squeeze the surplus of the product. Mop the floor with the wet mop.

Make sure to wet the surface completely. Allow to act for 5 minutes. Let it air dry. No rinsing needed after disinfection.

4.1.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.1.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.1.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.1.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.2. Use description

Table 2.

Use # 2.2 – Cleaning and disinfection with a cloth or sponge (incl. healthcare areas)

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data

Field(s) of use	indoor use Cleaning and disinfection product for hard surfaces (incl. healthcare areas).
Application method(s)	Method: spreading Detailed description: Spreading with a soaked cloth or sponge.
Application rate(s) and frequency	Application rate: 1 l to 40 ml/m ² Ready to use product Number and timing of application: Use at room temperature Frequency of use: 1 time a day
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	Professional and non-professional users: High Density Polyethene (HDPE) Canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres

4.2.1. Use-specific instructions

Soak the cloth or sponge in the product by dipping. Distribute the liquid on the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

4.2.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.2.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.2.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.2.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.3. Use description

Table 3.

Use # 2.3 – Cleaning and disinfection of small objects by soaking or dipping (incl. healthcare areas)

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Cleaning and disinfection product for small objects (e.g. scissors, brushes.) incl. objects for healthcare areas. The biocidal use excludes disinfection of medical devices.
Application method(s)	Method: Soaking or dipping Detailed description: Soaking or dipping in the solution.
Application rate(s) and frequency	Application rate: up to 3 litres Ready to use product Number and timing of application: Use at room temperature Frequency of use: 1 time a day
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	Professional and non-professional users: High Density Polyethene (HDPE)Canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres

4.3.1. Use-specific instructions

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

The solutions for soaking or dipping can be used only once.

4.3.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 2

4.3.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 2

4.3.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 2

4.3.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 2

4.4. **Use description**

Table 4.

Use # 2.4 – Cleaning and disinfection by foaming (incl. healthcare areas)

Product type	PT02: Disinfectants and algacides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Ready to use cleaning and disinfection product for hard surfaces (incl. healthcare areas).</p>
Application method(s)	<p>Method: spraying</p> <p>Detailed description: Spraying using a trigger foam. Optionally wipe to spread the solution.</p>
Application rate(s) and frequency	<p>Application rate: 1 l/m² Ready to use product</p> <p>Number and timing of application: Use at room temperature</p> <p>Frequency of use: For use in toilets and bathrooms: 4 times a day For use for cosmetic/pharmaceutical manufacturing facilities: 8 times a day For use in laboratories: 10 times a day For use in cleanrooms: 10 times a day</p>

	For use on a counter in a shop/supermarket: 8 times a day For use in hospitals and medical practices: 8 times a day
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	Professional and non-professional users: 750 ml High Density Polyethene (HDPE) trigger spray with foam head HDPE Canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres

4.4.1. Use-specific instructions

Apply the product on the surface. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surface disinfection.

4.4.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.4.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.4.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.4.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.5. Use description

Table 5.

Use # 2.5 – Cleaning and disinfection using moist wipes (incl. healthcare areas).

Product type	PT02: Disinfectants and algacides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data

	Scientific name: no data Common name: Viruses Development stage: no data
Field(s) of use	indoor use Ready to use cleaning and disinfection product for hard surfaces (incl. healthcare areas).
Application method(s)	Method: Wiping Detailed description: Wiping with prepared moist wipes.
Application rate(s) and frequency	Application rate: 1,5 m ² per wipe Ready to use product Number and timing of application: Frequency of use: 8 times a day Use at room temperature
Category(ies) of users	professional
Pack sizes and packaging material	HDPE Canister 1L - 1,5 L - 2 L - 2,5 L - 3 L - 5 L - 10 L - 20 L - 50 L - 100 L - 200 L HDPE IBC 100 - 1 000 Liter Dry Wipes are packed in a HDPE bucket with 125 wipes (PP) sold separately.

4.5.1. Use-specific instructions

Apply 1,5 litres product to the Dry Wipes bucket. Distribute the liquid evenly. Close the bucket.

Open the lid. Take a single wipe. Close the lid after taking a wipe. Only use wet wipes. Wipe the surface and make sure to wet the surface completely.

Allow to act for 5 minutes. Let it air dry. No rinsing needed after disinfection.

4.5.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.5.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.5.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.5.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.6. Use description

Table 6.

Use # 2.6 – Cleaning and disinfection by mopping (incl. food contact surfaces)

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Ready to use cleaning and disinfection product (incl. food contact surfaces)</p>
Application method(s)	<p>Method: Mopping</p> <p>Detailed description: Mopping with a soaked mop.</p>
Application rate(s) and frequency	<p>Application rate: 11 to 40 ml/m² Ready to use product</p> <p>Number and timing of application: Use at room temperature</p> <p>Frequency of use: 1 time a day</p>
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	<p>Professional and non-professional users: High Density Polyethene (HDPE) canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres</p> <p>Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres</p>

4.6.1. Use-specific instructions

Fill a container with the product. Dip the mop into the container. Squeeze the surplus solution. Mop the floor with the wet mop.

Make sure to wet the surface completely. Allow to act for 5 minutes and let it air dry. No rinsing needed after disinfection.

4.6.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.6.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 2

4.6.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 2

4.6.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 2

4.7. Use description

Table 7.

Use # 2.7 – Cleaning and disinfection with a cloth or sponge (incl. food contact surfaces)

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Ready to use cleaning and disinfection product for hard surfaces (incl. food contact surfaces).</p>
Application method(s)	<p>Method: spreading</p> <p>Detailed description: Spreading with a soaked cloth or sponge.</p>
Application rate(s) and frequency	<p>Application rate: 1 l to 40 ml/m² Ready to use product</p> <p>Number and timing of application: Use at room temperature Frequency of use: 1 time a day</p>

Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	Professional and non-professional users: High Density Polyethene (HDPE) canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres

4.7.1. Use-specific instructions

Soak the cloth or sponge in the product by dipping. Distribute the liquid on the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

4.7.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.7.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.7.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.7.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.8. Use description

Table 8.

Use # 2.8 – Cleaning and disinfection by soaking or dipping of small objects (incl. food contact surfaces)

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data

Field(s) of use	indoor use Ready to use cleaning and disinfection product for small objects (e.g. kitchen utensils) incl. food contact surfaces).
Application method(s)	Method: Soaking or dipping Detailed description: Soaking or dipping in the product.
Application rate(s) and frequency	Application rate: up to 3 litres Ready to use product Number and timing of application: Use at room temperature Frequency of use: 1 time a day
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	Professional and non-professional users: High Density Polyethene (HDPE) canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres

4.8.1. Use-specific instructions

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection. The solutions for soaking or dipping can be used only once.

4.8.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.8.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.8.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.8.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.9. Use description

Table 9.

Use # 2.9 – Cleaning and disinfection by foaming (incl. food contact surfaces)

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Ready to use cleaning and disinfection product for hard surfaces (incl. food contact surfaces).</p>
Application method(s)	<p>Method: spraying</p> <p>Detailed description: Spraying using a trigger foam. Optionally wipe to spread out the solution.</p>
Application rate(s) and frequency	<p>Application rate: 1 l/m² Ready to use product</p> <p>Number and timing of application: Use at room temperature</p> <p>Frequency of use for: For use in kitchens and canteen: 4 times a day For use in food processing industries: 4 times a day For use on a counter in a shop/supermarket: 8 times a day</p>
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	<p>Professional and non-professional users: 750 ml High Density Polyethene (HDPE) trigger foam HDPE Canister</p> <p>1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres</p> <p>Professional users: HDPE canister 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres</p>

4.9.1. Use-specific instructions

Apply the product on the surface. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Wipe dry or let it air dry. No rinsing needed after disinfection.

Only for small surface disinfection.

4.9.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.9.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.9.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.9.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.10. Use description

Table 10.

Use # 2.10 – Cleaning and disinfection with moist wipes (incl. food contact surfaces)

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Ready to use cleaning and disinfection product for hard surfaces (incl. food contact surfaces).</p>
Application method(s)	<p>Method: Wiping</p> <p>Detailed description: Wiping with prepared moist wipes.</p>
Application rate(s) and frequency	<p>Application rate: 1,5 m² per wipe Ready to use product</p> <p>Number and timing of application: Use at room temperature Frequency of use: 8 times a day</p>

Category(ies) of users	professional
Pack sizes and packaging material	RTU liquid: High Density Polyethene (HDPE) Canister 1 litres - 1,5 litres – 2 litres - 2,5 litres – 3 litres – 5 litres – 10 litres - 20 litres – 50 litres – 100 litres – 200 litres HDPE IBC 100 - 1 000 litres Dry Wipes are packed in a HDPE bucket with 125 wipes (PP). (sold separately)

4.10.1. Use-specific instructions

Apply 1,5 litres product to the Dry Wipes bucket. Distribute the liquid evenly. Close the bucket.

Open the lid, take a single wipe, close the lid after taking a wipe. Only use wet wipes. Wipe the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Let it air dry. No rinsing needed after disinfection.

4.10.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 2

4.10.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 2

4.10.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 2

4.10.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of Meta-SPC 2

4.11. Use description

Table 11.

Use # 2.11 – Disinfection of pre-cleaned hard surfaces

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: no data Common name: Bacteria Development stage: no data Scientific name: no data Common name: Yeasts Development stage: no data Scientific name: no data Common name: Viruses Development stage: no data

Field(s) of use	indoor use Disinfection product for pre-cleaned hard surfaces (incl. healthcare areas, pharmaceutical industry, cosmetic industry, laboratories and cleanrooms)
Application method(s)	Method: Spreading, dipping and foam spraying Detailed description: Spreading with a soaked mop. Spreading with a soaked cloth or sponge. Dipping in a solution. Foam spraying, optionally wipe to spread the solution.
Application rate(s) and frequency	Application rate: Application using a soaked mop: 11 to 40 ml/m ² , application using a soaked cloth/sponge: 11 to 40 ml/m ² , dipping: 3 litres, foam spraying: 11 ml/m ² ready to use product Number and timing of application: Frequency of use: Application using a soaked mop: 7 times a day Application using a soaked cloth/sponge: 7 times a day Dipping: 1 time a day Foam spraying: bathrooms & toilets 4 times a day healthcare 8 times a day laboratories & cleanrooms 10 times a day Use at room temperature
Category(ies) of users	professional
Pack sizes and packaging material	High Density Polyethene (HDPE) canister: 1 litres - 1,5 litres - 2 litres - 2,5 litres - 3 litres - 5 litres - 10 litres - 20 litres - 50 litres - 100 litres - 200 litres HDPE bottle (J-Flex): 1,5 litres HDPE Bottle (Smart Dose): 1,4 litres - 1,5 litres HDPE IBC: 100-1 000 litres

4.11.1. Use-specific instructions

Contact time: 5 minutes

Mopping: Dip the mop into the container. Squeeze the surplus solution. Apply onto floor.

Make sure to wet the surface completely. Allow to act for the required contact time.

Cloth/sponge application: Soak the cloth or sponge in the solution. Distribute the liquid on the surface.

Make sure to wet the surface completely. Allow to act for the required contact time.

Dipping:

Cover dipping bath with a lid after placing the objects in the disinfection solution.

Ensure that items to be disinfected are fully submerged in the solution for 5 minutes.

The solutions for soaking or dipping can be used only once.

Foam spraying: Apply the product. Optionally wipe to spread the solution.

Make sure to wet the surface completely. Allow to act for the required contact time.

Only for small surfaces disinfection.

Wipe dry or let air dry. No rinsing needed after disinfection.

4.11.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 2

4.11.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 2

4.11.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 2

4.11.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 2

5. **GENERAL DIRECTIONS FOR USE OF THE META SPC 2**

5.1. **Instructions for use**

For professional use:

Keep away from food, drink and animal feeding stuffs.

Wash hands after handling.

Keep only in original container.

Do not mix with other products.

See use-specific instructions for use.

For non-professional use:

Comply with the instructions for use.

Keep away from food, drink and animal feeding stuffs.

Wash hands after handling.

Keep only in original container.

Do not mix with other products.

See use-specific instructions for use.

5.2. **Risk mitigation measures**

Keep pets away from treated surfaces until dried.

5.3. **Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

IF INHALED: If symptoms occur call a POISON CENTRE or a doctor.

IF SWALLOWED: If symptoms occur call a POISON CENTRE or a doctor.

IF ON SKIN: Wash skin with water. If symptoms occur call a POISON CENTRE or a doctor.

IF IN EYES: If symptoms occur rinse with water. Remove contact lenses, if present and easy to do. Call a POISON CENTRE or a doctor..

If medical advice is needed, have product container or label at hand.

5.4. Instructions for safe disposal of the product and its packaging

Dispose of contents/container to a hazardous waste disposal service in accordance with the statutory regulations.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

2 years shelf life.

Store in the dark. Do not store at temperatures above 50°C.

Keep out of reach of children and non-target animals/pets.

6. OTHER INFORMATION

The products of this meta-SPC may also be intended for the disinfection of medical devices' surfaces in medical and clinical areas. In such a case, their placing on the market or putting into service must comply with Regulation (EU) 2017/745 of the European Parliament and of the Council of 5 April 2017 on medical devices, amending Directive 2001/83/EC, Regulation (EC) No 178/2002 and Regulation (EC) No 1223/2009 and repealing Council Directives 90/385/EEC and 93/42/EEC (OJ L 117, 05/05/2017, p. 1) and they are medical devices classified as class IIa. Those uses are out of the scope of this biocidal product family authorisation and are not covered by it.

With respect to the "Category(ies) of users" note: "Professionals (including industrial users) means trained professionals if this is required by national legislation.

7. THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 2

7.1. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)	Oxivir Excel Foam	Market area: EU
	Oxivir Excel Foam H+	Market area: EU
	Oxivir Excel Spray	Market area: EU
	Oxivir Excel Spray H+	Market area: EU
	Oxivir Excel RTU	Market area: EU
	Oxivir Excel RTU H+	Market area: EU
	Oxivir Excel RTU IBC	Market area: EU

	Oxivir Excel Foam Acti	Market area: EU
	Oxivir Excel Foam Des	Market area: EU
	Oxivir Excel Foam Duo	Market area: EU
	Clinell H2O2 RTU	Market area: EU
	Antibac Oxivir RTU	Market area: EU
	Divosan AHP Foam RTU VS71	Market area: EU
	TASKI Sani AHP- Des Spray	Market area: EU
	TASKI Sprint AHP-Des Spray	Market area: EU
	Suma AHP RTU D10.7	Market area: EU
	Suma AHP Spray D10.7	Market area: EU
	Suma AHP RTU	Market area: EU
	Suma AHP Spray	Market area: EU
	ClearKlens AHP RTU	Market area: EU

	ClearKlens Excel RTU	Market area: EU			
	DOMES- TOS Expert	Market area: EU			
	GLORIX Expert	Market area: EU			
	KLINEX Expert	Market area: EU			
	LYSO- FORM Expert	Market area: EU			
	SAVO Expert	Market area: EU			
	CIF Expert	Market area: EU			
	VISS Expert	Market area: EU			
	JIF Expert	Market area: EU			
	FLORAS- ZEPT Expert	Market area: EU			
	ANDY Expert	Market area: EU			
	SUPER- ANDY Expert	Market area: EU			
	LIFEBUOY Expert	Market area: EU			
	ANDY- SAN Expert	Market area: EU			
Authorisation number		EU-0032483-0002 1-2			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Hydrogen peroxide		active substance	7722-84-1	231-765-0	0,36 % (w/w)

1. META SPC 3 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 3 identifier

Identifier	Meta SPC: Meta-SPC 3: Wipes
------------	-----------------------------

1.2. Suffix to the authorisation number

Number	1-3
--------	-----

1.3. Product type(s)

Product type(s)	PT02: Disinfectants and algaecides not intended for direct application to humans or animals PT04: Food and feed area
-----------------	---

2. META SPC 3 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 3

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Hydrogen peroxide		active substance	7722-84-1	231-765-0	0,36 - 0,36 % (w/w)

2.2. Type(s) of formulation of the meta SPC 3

Formulation type(s)	AL (Any other liquid) - RTU wipe
---------------------	----------------------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 3

Hazard statements	H290: May be corrosive to metals.
Precautionary statements	P102: Keep out of reach of children. P234: Keep only in original packaging.

4. AUTHORISED USE(S) OF THE META SPC

4.1. Use description

Table 1.

Use # 3.1 – Cleaning and disinfection using moist wipes on hard surfaces incl. healthcare areas

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Wipe for cleaning-disinfecting for hard surfaces incl. healthcare areas.</p>
Application method(s)	<p>Method: Wiping</p> <p>Detailed description: Wiping with ready to use moist wipes.</p>
Application rate(s) and frequency	<p>Application rate: 1,5 m² per wipe Ready to use wipe.</p> <p>Number and timing of application: Frequency of use: 8 times a day Use at room temperature</p>
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) bucket or tub with 100 PP wipes Polyethylene terephthalate (PET)/low-density polyethylene (LDPE) flow-pack with 100 polypropylene (PP) wipes PET/LDPE standing bag with 100 PP wipes All packages are both for professional and non-professional use</p>

4.1.1. Use-specific instructions

Open the lid. Take a single wipe. Close the lid after taking a wipe. Only use wet wipes. Wipe the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Let it air dry. No rinsing needed after disinfection.

4.1.2. Use-specific risk mitigation measures

See general directions for use of Meta-SPC 3

4.1.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of Meta-SPC 3

4.1.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of Meta-SPC 3

4.1.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 3

4.2. **Use description**

Table 2.

Use # 3.2 – Cleaning and disinfection using moist wipes on hard surfaces incl. food contact surfaces

Product type	PT04: Food and feed area
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Wipe for cleaning-disinfecting hard surfaces (incl. food contact surfaces)</p>
Application method(s)	<p>Method: Wiping</p> <p>Detailed description: Wiping with ready to use moist wipes.</p>
Application rate(s) and frequency	<p>Application rate: 1,5 m² per wipe Ready to use wipe.</p> <p>Number and timing of application: Use at room temperature Frequency of use: 8 times a day</p>
Category(ies) of users	professional ; general public (non-professional)
Pack sizes and packaging material	<p>High Density Polyethene (HDPE) bucket or tub with 100 PP wipes PET or LDPE flow-pack with 100 PP wipes PET or LDPE standing bag with 100 PP wipes All packages are both for professional and non-professional use</p>

4.2.1. *Use-specific instructions*

Open the lid. Take a single wipe. Close the lid after taking a wipe. Only use wet wipes. Wipe the surface.

Make sure to wet the surface completely. Allow to act for 5 minutes.

Let it air dry. No rinsing needed after disinfection.

4.2.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 3

4.2.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 3

4.2.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 3

4.2.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 3

4.3. **Use description**

Table 3.

Use # 3.3 – Wipe for disinfecting pre-cleaned hard surfaces

Product type	PT02: Disinfectants and algaecides not intended for direct application to humans or animals
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	<p>Scientific name: no data Common name: Bacteria Development stage: no data</p> <p>Scientific name: no data Common name: Yeasts Development stage: no data</p> <p>Scientific name: no data Common name: Viruses Development stage: no data</p>
Field(s) of use	<p>indoor use</p> <p>Disinfection wipe for pre-cleaned hard surfaces (incl. healthcare areas, pharmaceutical industry, cosmetic industry, laboratories and cleanrooms)</p>
Application method(s)	<p>Method: Wiping</p> <p>Detailed description: Wiping with ready to use moist wipes.</p>
Application rate(s) and frequency	<p>Application rate: 1,5 m² per wipe Ready to use</p> <p>Number and timing of application: Use at room temperature Frequency of use: 8 times a day</p>
Category(ies) of users	professional

Pack sizes and packaging material	HDPE bucket/tub with 100 PP wipes PET/LDPE flow-pack with 100 PP wipes PET/LDPE standing bag with 100 PP wipes
-----------------------------------	--

4.3.1. *Use-specific instructions*

Contact time: 5 minutes

Open the lid. Take a single wipe. Close the lid after taking a wipe. Only use wet wipes. Wipe the surface.

Make sure to wet the surface completely. Allow to act for the required contact time.

Let it air dry. No rinsing needed after disinfection.

4.3.2. *Use-specific risk mitigation measures*

See general directions for use of Meta-SPC 3

4.3.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

See general directions for use of Meta-SPC 3

4.3.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

See general directions for use of Meta-SPC 3

4.3.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

See general directions for use of Meta-SPC 3

5. **GENERAL DIRECTIONS FOR USE OF THE META SPC 3**

5.1. **Instructions for use**

For professional use:

Keep away from food, drink and animal feeding stuffs.

Wash hands after handling.

Keep only in original container.

Do not mix with other products.

See use-specific instructions for use.

For non-professional use:

Comply with the instructions for use.

Wash hands after handling.

Keep only in original container.

Do not mix with other products.

See use-specific instructions for use.

5.2. **Risk mitigation measures**

Keep pets away from treated surfaces until dried.

5.3. **Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

IF INHALED: If symptoms occur call a POISON CENTRE or a doctor.

IF SWALLOWED: If symptoms occur call a POISON CENTRE or a doctor.

IF ON SKIN: Wash skin with water. If symptoms occur call a POISON CENTRE or a doctor.

IF IN EYES: If symptoms occur rinse with water. Remove contact lenses, if present and easy to do. Call a POISON CENTRE or a doctor.

If medical advice is needed, have product container or label at hand.

5.4. **Instructions for safe disposal of the product and its packaging**

Dispose of contents/container to a hazardous waste disposal service in accordance with the statutory regulations.

5.5. **Conditions of storage and shelf-life of the product under normal conditions of storage**

2 years shelf life.

Store in the dark. Do not store at temperatures above 50°C.

Keep out of reach of children and non-target animals/pets.

6. **OTHER INFORMATION**

The products of this meta-SPC may also be intended for the disinfection of medical devices' surfaces in medical and clinical areas. In such a case, their placing on the market or putting into service must comply with Regulation (EU) 2017/745 of the European Parliament and of the Council of 5 April 2017 on medical devices, amending Directive 2001/83/EC, Regulation (EC) No 178/2002 and Regulation (EC) No 1223/2009 and repealing Council Directives 90/385/EEC and 93/42/EEC (OJ L 117, 05/05/2017, p. 1) and they are medical devices classified as class IIa. Those uses are out of the scope of this biocidal product family authorisation and are not covered by it.

With respect to the "Category(ies) of users" note: "Professionals (including industrial users) means trained professionals if this is required by national legislation.

The following types of wipes are included:

Wipe with size 15 x 25 cm; weight is 1,52 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 15,2 x 17,7 cm; weight is 1,51 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 15,2 x 20,3 cm; weight is 1,51 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 17,7 x 20,3 cm; weight is 1,50 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 19 x 25,4 cm; weight is 1,49 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 20 x 26,7 cm; weight is 1,49 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 20 x 30 cm; weight is 1,50 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 25 x 25 cm; weight is 1,51 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 25,4 x 25,4 cm; weight is 1,51 gr; 3,75 ml liquid per wipe (wipe liquid load is 250%)

Wipe with size 15 x 25 cm; weight is 1,52 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 15,2 x 17,7 cm; weight is 1,51 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 15,2 x 20,3 cm; weight is 1,51 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 17,7 x 20,3 cm; weight is 1,50 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 19 x 25,4 cm; weight is 1,49 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 20 x 26,7 cm; weight is 1,49 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 20 x 30 cm; weight is 1,50 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 25 x 25 cm; weight is 1,51 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

Wipe with size 25,4 x 25,4 cm; weight is 1,51 gr; 6 ml liquid per wipe (wipe liquid load is 400%)

7. THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 3

7.1. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)	Oxivir Excel Wipe	Market area: EU
	Oxivir Excel Wipe H+	Market area: EU
	Oxivir Excel Wipe Acti	Market area: EU
	Oxivir Excel Wipe Des	Market area: EU
	Oxivir Excel Wipe Duo	Market area: EU
	Clinell H2O2 wipes	Market area: EU
	Antibac Oxivir wipes	Market area: EU
	Divosan AHP Wipes VS72	Market area: EU
	Suma AHP Wipes	Market area: EU
	Suma San AHP Wipes	Market area: EU

	ClearKlens AHP Wipes	Market area: EU
	ClearKlens Excel Wipes	Market area: EU
	DOMES- TOS Expert Wipes	Market area: EU
	GLORIX Expert Wipes	Market area: EU
	KLINEX Expert Wipes	Market area: EU
	LYSO- FORM Expert Wipes	Market area: EU
	SAVO Expert Wipes	Market area: EU
	CIF Expert Wipes	Market area: EU
	VISS Expert Wipes	Market area: EU
	JIF Expert Wipes	Market area: EU
	FLORAS- ZEPT Expert Wipes	Market area: EU
	ANDY Expert Wipes	Market area: EU
	SUPER- ANDY Expert Wipes	Market area: EU

		LIFEBUOY Expert Wipes	Market area: EU		
		ANDYSAN Expert Wipes	Market area: EU		
Authorisation number		EU-0032483-0003 1-3			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Hydrogen peroxide		active substance	7722-84-1	231-765-0	0,36 % (w/w)